



**Albac®**  
*Poultry*

## PRODUCT DESCRIPTION

Albac® contains feed grade zinc bacitracin equivalent to 150 grams (15%) bacitracin per kilogram.

## INDICATIONS FOR USE

1. Supporting the prevention and control of necrotic enteritis in chickens caused or complicated by *Clostridium spp.*
2. Increased weight gain and improved feed efficiency.
3. Increased egg production and improved feed efficiency in laying hens.

## KEY POINTS

- Albac® helps lowering intestinal counts of *C. perfringens* as well as *Lactobacillus salivarius*, this promotes a healthy gut flora and higher nutrient absorption in broilers.
- The only bacitracin available for use in all ages of laying hens.
- Not absorbed from the intestinal tract, thus no withdrawal period is required prior to slaughter.
- Consistent performance after long time use.
- Patented Chilsonation process improves flowability and stability, and minimizes dust to ensure proper dosage.
- Heat stress does not affect the performance of zinc bacitracin.

## USERS INFORMATION: ALBAC® 15%

| <b>Active Drug Ingredient</b>                  | <b>Each kilogram contains feed grade zinc bacitracin equivalent to 150 g bacitracin (Master Standard).</b>                                                                                          |                                  |                                                                                                                                                                                                |
|------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Composition</b>                             | <b>The zinc salt of a dried fermentation product obtained by culturing <i>Bacillus licheniformis</i> Tracy on media adapted for microbiological production of bacitracin and calcium carbonate.</b> |                                  |                                                                                                                                                                                                |
| <b>For Use in Manufactured Medicated Feeds</b> |                                                                                                                                                                                                     |                                  |                                                                                                                                                                                                |
| Species                                        | Grams bacitracin per tonne of Feed                                                                                                                                                                  | Albac® dosage                    | Indications for Use                                                                                                                                                                            |
| <b>Broiler chickens</b>                        | 5 – 50                                                                                                                                                                                              | dose between 35 - 335 gram /mton | For increased weight gain and improved feed efficiency                                                                                                                                         |
|                                                | 50                                                                                                                                                                                                  | dose of 335 gram/mton            | Supporting the prevention of necrotic enteritis. Medicated feed should be fed to poultry during the period when they are at most risk of the disease, which is usually between 2 and 12 weeks. |
|                                                | 200                                                                                                                                                                                                 | dose of 1335 gram/mton           | Supporting the treatment of necrotic enteritis: add Albac® for 5 – 7 days.                                                                                                                     |
| <b>Laying hens</b>                             | 15 – 100                                                                                                                                                                                            | dose between 100-667 gram/mton   | For increased weight gain and improved feed efficiency                                                                                                                                         |

Can be used in combination with our registered feed additives and coccidiostats\*.

## HANDLING

- Albac® is packaged in a 25 kg, multilayer paper bag with protective barrier ply.
- Store in a cool (<25°C), dry place and keep package closed to avoid contamination.
- Expiry data is 36 months from the date of manufacture.
- For use in dry feeds only
- Not for use in feeds for breeder chickens

\*Specific claims vary by country.

Please contact your Huvepharma representative or distributor for more information.

References available on request.

