



Enrofloxacin 10%

ENROPHARMA®

Antibacterial / Oral Solution



Each Liter of product contains:

Enrofloxacin.....100 g
Excipient q.s. to.....1 Liter

Target Species: Poultry

Mode of action:

Enrofloxacin is a synthetic broad spectrum antibiotic and bactericidal belongs to the Quinolone group that it inhibits the activity of DNA-gyrase enzyme. The viability of susceptible organisms to Enrofloxacin is usually lost within 20-30 min of exposure to optimal antibacterial concentration, and effective against a wide range of gram negative and gram positive bacteria as well as mycoplasma.

Indications:

ENROPHARMA® is being used as an aid in control and treatment diseases of the respiratory and alimentary tracts, urinary-reproductive organs of poultry caused by gram-positive and gram-negative bacteria or mycoplasma origin such as: *CRD, Mycoplasmosis, Pasteurellosis (Fowl cholera), Fowl typhoid, Paratyphoid, E. coli infection, Colibacillosis, Actinobacillosis, Salmonellosis, Brucellosis, Coli septicemia, Staphylococcal and Streptococcal infection*, in secondary bacterial infections following viral infections and etc. it is used by drinking water.

Contradictions:

Do not use in case of hypersensitivity to Enrofloxacin.
Do not use in layers producing eggs for human consumption.

Dosage and mode of administration:

10 mg per kg of body weight per day (equivalent to 500 ml of ENROPHARMA® per 1000 liters of drinking water) for 3 to 5 Consecutive days.

Notice:

Duration and amount of use, should be administered by veterinarian.
Medicated solution should be freshly prepared daily and consumed within 24 hours.

Medicated drinking water should be the only water source available.

After opening the bottle, the contents should be consumed within 30 days.

The dosage rates of oral solutions or water soluble powder depending on the amount of up take of medicated drinking can be influenced by factors like ambient temperature (in different seasons hot or cold climate), salt content of food, clinical conditions of animal, stocking density, quality and hardness of water.

Interactions: Should not be used with chloramphenicol, macrolides, tetracycline and aminoglycosides. Magnesium and Aluminum containing substances might hamper the absorption of Enrofloxacin from Gastro-Intestinal tract.

Withdrawal Time: 8 days before slaughtering.

Precaution: For veterinary use only. Keep out of reach of children. Wear overalls, impervious gloves, safety glasses and dust mask when handling or mixing the product. If ENROPHARMA® accidentally enters the eyes, wash eyes with plenty amount of water.

Storage: Keep in closed original pack, cool and dry, 10- 25°C. Protect from sunlight

Shelf life: 2 years from the date of manufacture.

Packing: 1 liter Polyethylene Bottles.

