



Clopidol 25%

CLOPIDAL®

Synthetic Coccidiostat / Premix



Each kg of product contains:

Clopidol 250 g
Excipient q.s. to 1000 g

Target Species: Poultry

Mode of action:

Clopidol is a broad-spectrum synthetic coccidiostat that prevents energy metabolism in the cytochrome oxidase system of coccidia. Clopidol leads to an inhibition of the development of the first generation of schizonts by effecting on the intracellular sporozoites.

Clopidol is effective against coccidiosis caused by different species *Eimeria* such as: *Eimeria tenella*, *E. acervulina*, *E. maxima*, *E. necatrix*, *E. brunette*, *E. mitis* and *E. praecox*.

Indications:

CLOPIDAL® is being used as an aid to prevent of coccidiosis in broiler chickens and chickens reared for laying up to 16 weeks ages.

Contraindications:

Do not use to laying hens for more than 16 weeks.

Dosage and mode of administration:

Broiler chickens and chickens reared for laying up to 16 weeks of age: 500 gram **CLOPIDAL®** per ton of feed (equivalent 125 ppm clopidol).

Notice:

Duration and amount of use should be administered by veterinarian. In order to reach uniform homogenization with feed, it is recommended to mix the measured quantity of the preparation (calculated on the basis of the prescribed dose) on stages in the following order up to 10 kg of feed, up to 100 kg of feed, and up to 1000 kg of feed.

Interactions: **CLOPIDAL®** should not be mixed with any other products that has the same effects.

Withdrawal Time: 5 days before slaughtering.

Precaution: For veterinary use only. Keep out of reach of children. Wear overalls, impervious gloves, safety glasses and dust mask when handling or mixing the product. If **CLOPIDAL®** accidentally enters the eyes wash eyes with plenty amount of water.

Storage: Keep in closed original pack, cool and dry place, below 25°C. Protect from direct sunlight.

Shelf life: 2 years from the date of manufacture.

Packing: 5 kg three-layers aluminum laminated foil.

