

Each Liter of product contains:

Vitamin A (Palmitate) 10,00	0,000 IU
Vitamin D3 (Crystalline cholecalciferol) 3,00	0,000 IU
Vitamin E (DI-a-Tocopherol Acetate)	5 g
Vitamin B1 (Thiamin Hydrochloride)	2g
Vitamin B2 (Riboflavin 5-Sodium phosphate)	2g
Dexpanthenol	5 g
Nicotinamide	10 g
Vitamin B6 (Pyridoxine Hydrochloride)	1g
Sodium Chloride	10 g
Potassium Chloride	1g
Potassium Acetate	2g
Magnesium Cloride	3g
Sorbitol	50 g
Excipient q.s. to	1lit

Target Species: Livestock and Poultry

Mode of action:

Vitamins are compounds that play a vital and unique role in all vital functions of the body, including maintaining tissues, resistance to pathogens and stress, enhancing the body's immunity and proper functioning of the organs. Since the body of animals is not able to make these compounds, they should be taken orally in their diet. Also, electrolytes are substances whose balance in the body is very important and their deficiencies can lead to various complications

Indications:

POLYLIT[®] by maintaining optimum hydration helps support digestion, feed intake and electrolyte status. It supplies vital electrolytes in any type of dehydration caused by diarrhea, during or after antibiotic therapy, parasitic and coccidiosis treatment. **POLYLIT**[®] is used as an aid to improve lesions caused by vitamins deficiency in livestock and poultry.

It improves the immune response in stress – related situations, such as: vaccination, weaning process, transportation, feed changes, debeaking etc.

Dosage and mode of administration:

Poultry: 1 liter per 800 to 1000 liters of drinking water for 3 to 5 days.

Lambs and Goats: 0.5 to 1 ml per animal for 3 days. Calves: 1 to 2 ml per animal for 3 days. Horse: 1 ml per 100 kg body weight for 3 days.

Notice:

Duration and amount of use, should be administered by veterinarian. Prepared solution should be consumed within 18 hours. After opening the bottle, the contents should be consumed within 20 days. Administration oral solution or water soluble powder in different seasons (hot or cold climate) depends on consumption of water by the target animal.

The gradual darkening of the product does not affect its stability. If stored at an inappropriate temperature, there is a possibility of precipitation of the product.

Withdrawal Time: None

Precaution: For veterinary use only. Keep out of reach of children. **Storage:** Keep in closed original pack, dry and dark place, 10-25°C.

Protect from direct sunlight.

Shelf life: 2 years from the date of manufacture. **Packing:** 1 liter Polyethylene Bottles.

