



500g

Vitamin C 50%

RUMITAMINE®

Vitamin/ Water Soluble Powder



Each kg of product contains:

Vit C (Ascorbic Acid)	500 g
Excipient q.s. to	1000 g

Target Species: Poultry

Mode of action:

Vitamin C plays an important physiologic role in numerous metabolic functions, including synthesis and production of hormones involved in resistance to stress, amelioration of oxidative stress and immune regulation.

Indications:

- Stress conditions due to vaccination, debeaking, transporting, relocating, changes of feed, weather and viral and parasitical diseases.
- Improves quality, fertility and hatchability rates, increases the egg production and hardness of the egg shell.
- Increases the body resistance against infections and the immune response.
- Facilitates absorption and the almost utilization of iron, calcium, selenium, copper and other nutrients.
- Reduces the toxic effects of fungi and other disease through its oxidizing – reducing and Anti-hemorrhagic action.

Dosage and mode of administration:

Heat stress, production stress in layers, stress due to relocating: 6-12 g in 100 lit of drinking water (10 -15 mg ascorbic acid per kg of live body weight) at least for 3 days.

Stress of vaccination or beak trimming, cold or heat stress: 12-20 g in 100 liter of drinking water (15-20 mg ascorbic acid per kg of live body weight) at least for 3 days.

Parasitic and infectious diseases: 20 g in 100 liter of drinking water (30-40 mg ascorbic acid per kg of live body weight) at least for 3 days.

Notice:

Prepared solution should be consumed within 12 hours. Administration oral solution or water soluble powder in different seasons (warm or cold weather) depends on consumption of water by the target animal.

Withdrawal Time: None

Storage: Keep in closed original pack, cool and dry place, below 25°C. Protect from direct sunlight.

Shelf life: 2 years from the date of manufacture.

Packing: 5 kg three-layered aluminum laminated foil.

