



Lincomycin 0.88%

AMIRAFEED[®] 8.8

Antibiotic / Premix



Each kg of product contains:

Lincomycin (as Hydrochloride) 8.8 g
Excipient q.s. to 1000 g

Target Species: Broiler Poultry

Mode of action:

Lincomycin is a bacteriostatic antibiotic from the lincosamides group that prevents the synthesis of bacterial proteins by binding to the 50S ribosomal subunit. And with the above mechanism, it is effective on gram-positive, bacteria such as; streptococci, staphylococci, clostridium and mycoplasma.

Indications:

It is used to control necrotic enteritis caused by Clostridium perfringens and other microorganisms, improving growth rate and food conversion rate in broilers and to control CRD.

Contraindications:

Do not use in Breeder, Layer poultry and turkeys.

Dosage and mode of administration:

For necrotic intestinal inflammation, CRD control and improvement of growth rate and improvement of feed conversion ratio: 500 grams per ton of feed (from the first week of raising broilers until the end of the period)

Notice:

Duration and amount of use should be administered by veterinarian. To get a uniform mix, at first should be mix the Lincomycin 8.8 premix based on the prescribed dosage, Respectively, with 10 kg feed, then, with 100 kg and finally, with 1000 kg feed. Mix Carefully to achieve homogeneous mixture.

Interactions: Simultaneous use with erythromycin and tylosin is not recommended. It is compatible with all type of coccidiostats and growth promoters.

Withdrawal Time: Non.

Precaution: For veterinary use only. Keep out of reach of children. Wear protective clothes, impervious gloves and dust mask during mixing with feed if accidentally enters the eyes wash eyes with plenty amount of water.

Storage: Keep in closed original pack, cool and dry place, below 25°C. Protect from direct sunlight.

Shelf life: 2 years from the date of manufacture.

Packing: 5 kg three-layers aluminum laminated foil.

